

Positive Measure Spectrum for Schrödinger Operators with Periodic Magnetic Fields

Michael J. Gruber*

February 7, 2008

Abstract

We study Schrödinger operators with periodic magnetic field in $\mathbb{R}^2$, in the case of irrational magnetic flux. Positive measure Cantor spectrum is generically expected in the presence of an electric potential. We show that, even without electric potential, the spectrum has positive measure if the magnetic field is a perturbation of a constant one.

Introduction

Magnetic Schrödinger operators have been studied in Solid State Physics, especially in connection with the Quantum Hall effect, as well as on their own right. In a regular crystal ‘physics’ is periodic, i.e., the electric potential - caused by the background field of the ions - is a periodic function. Magnetic fields - internal as well as external ones - are periodic as well, the latter ones typically being constant. Alas, as is well known, magnetic fields enter the Schrödinger operator through a vector potential, so that the resulting operator is not necessarily periodic. Indeed, it is so only in the simple and well-understood case of ‘zero flux’, where one has absolutely continuous spectrum and band-structure (Birman & Suslina, 1998; Sobolev, 1999). Here, the magnetic flux (in units of flux quanta) is defined by

$$\Phi = \frac{1}{2\pi} \int_{\mathcal{F}} B(x, y) dx dy, \quad (1)$$

*Department of Mathematics, The University of Arizona, Tucson AZ, USA

where B is the magnetic field and $\mathcal{F}$ is a lattice cell (fundamental domain for the action of the group). Note that in our units with $\hbar = e = 1$, the magnetic flux quantum is just 2π .

For integer or rational flux the spectrum will still consist of bands (possibly degenerating into points), but pure point spectrum is possible.

For irrational flux one expects Cantor spectrum (i.e. a nowhere dense set, no isolated points). The question is now: If B is constant, what ‘defines the lattice’, thus defining $\mathcal{F}$ and the flux? A potential $V(x, y) = \cos(2\pi x) + \cos(2\pi y)$ has periods $(1, 1)$; one might consider any pair (m, n) of integers as periods of V , i.e., one can consider any coarser superlattice of $\mathbb{Z} \times \mathbb{Z}$ as the lattice of symmetry, but no finer lattice. Indeed, in this case one finds Cantor structure for a certain set of irrational values of Φ (Helffer & Sjöstrand, 1989).

On the other hand, $V(x, y) = \cos(2\pi x)$ has periods $(1, c)$ for any real c ; it does not define a fixed ‘minimally coarse’ lattice. Indeed, the Schrödinger operator with constant magnetic field and this potential V has band spectrum. This is still true for every potential $V(x, y) = V_0(x)$ with reasonably non-degenerate V_0 . Now, if we perturb such a V by a periodic V_1 we expect Cantor spectrum, although this is not known. We only know that the Schrödinger operator can be approximated in norm resolvent sense by operators from a natural algebra which have Cantor spectra (Gruber, 2001).

Setting the Cantor issue aside, Dinaburg et al. (1997) showed that what survives under this kind of perturbation is the positive measure of the bands, although the bands might dissolve into a Cantor set.

We ask the same question for Schrödinger operators with periodic magnetic field B , without electric potential V . For constant B the spectrum is pure-point and infinitely degenerate (Landau levels). Is a periodic (zero-flux) perturbation of B enough to cause the same effects as the potential V ?

In the course of giving an affirmative answer, we construct the generalized eigenfunctions and give estimates on the measure of the ‘bands’.

Outline

In Section 1 we describe the setup and perform a first perturbation by one-dimensional magnetic potentials. For the general case, we give the direct integral decomposition and express the operator in an appropriate basis ‘moving along the fiber’. In Section 2 we give concise formulations of the main results. The resulting double-infinite matrix problem is reduced in Section 3 to a problem which is almost diagonal in a sense made precise there. In

Section 4 we prove the main reduction lemma, using estimates on Weber-Hermite functions. Section 5 finishes the proof of the main results.

Acknowledgements

We thank Leonid Friedlander for fruitful discussions and exchange of ideas.

1 Setup and first perturbation

For simplicity of the notation we assume that the lattice is $\mathbb{Z} \times \mathbb{Z}$. Let B be an arbitrary smooth periodic magnetic field and Φ its flux through a fundamental cell of the lattice in units of flux quanta. Then we can decompose B as

$$\begin{aligned} B &= B_c + B_z \quad \text{with} \\ B_c &= 2\pi\Phi \quad \text{and} \\ B_z &= B - B_c \end{aligned}$$

Note that B_z has zero flux! Therefore, B_z has a periodic vector potential A_z , and we can choose it to be of the form

$$A_z(x, y) = \begin{pmatrix} \varepsilon_0 A^0(y) \\ \varepsilon_1 A^1(x, y) \end{pmatrix} \quad (2)$$

with periodic smooth A^0 and A^1 , where we introduced parameters $\varepsilon_0, \varepsilon_1$ for later convenience.

B_c is obviously constant, and we choose a vector potential

$$A_z(y) = B_c \begin{pmatrix} y \\ 0 \end{pmatrix} \quad (3)$$

for it. In this gauge, the magnetic Schrödinger operator takes on the form

$$H = \frac{1}{2} \left[\left(\frac{1}{i} \frac{\partial}{\partial x} - B_c y - \varepsilon_0 A^0(y) \right)^2 + \left(\frac{1}{i} \frac{\partial}{\partial y} - \varepsilon_1 A^1(x, y) \right)^2 \right]. \quad (4)$$

Note that, in order to emphasize the dependencies, we write out the arguments of A^0 and A^1 even though they are multiplication operators; in the

following we will keep sacrificing strictness of notation for better clarity in cases such as this one.

First we will set $\varepsilon_1 = 0$ and study the effect of switching on ε_0 . The operator (4) does not depend on x if $\varepsilon_1 = 0$. Using Fourier transform on $L^2(\mathbb{R}_x)$, we can decompose $L^2(\mathbb{R}^2)$ into a direct integral $\int_{\mathbb{R}}^{\oplus} L^2(\mathbb{R}_y) d\xi$ such that

$$L^2(\mathbb{R}^2) \ni f \mapsto \hat{f}, \quad \hat{f}_{\xi}(y) = \int_{\mathbb{R}} f(x, y) e^{-2\pi i \xi x} dx, \quad (5a)$$

$$f(x, y) = \int_{\mathbb{R}} \hat{f}_{\xi}(y) e^{2\pi i \xi x} d\xi; \quad (5b)$$

$$H = \int_{\mathbb{R}}^{\oplus} \hat{H}_{\xi} d\xi, \quad \hat{H}_{\xi} = V_{\xi}(y) - \frac{1}{2} \frac{d^2}{dy^2} \quad \text{with} \quad (5c)$$

$$V_{\xi}(y) = \frac{1}{2} (2\pi\xi - B_c y - \varepsilon_0 A^0(y))^2. \quad (5d)$$

In the case $\varepsilon_0 = \varepsilon_1 = 0$ we are dealing with the Landau Hamiltonian. We will go through its analysis since we will use its eigenfunctions later on as a basis.

1.1 $\varepsilon_0 = 0 = \varepsilon_1$

$$V_{\xi}(y) = \frac{1}{2} (2\pi\xi - B_c y)^2 = \frac{B_c^2}{2} (y - \beta\xi)^2 \quad (6)$$

is a harmonic oscillator potential with frequency B_c , shifted by $\beta\xi$ with $\beta = \frac{2\pi}{B_c} = \frac{1}{\Phi}$ (we assume $B_c \neq 0$, or else there is not much to do). If we denote by

$$\Omega_m(y) = \frac{(-1)^m}{\sqrt{\sqrt{\pi} 2^m m!}} \exp\left(\frac{y^2}{2}\right) \frac{d^m}{dy^m} \exp(-y^2), \quad m \in \mathbb{Z}_+, \quad (7)$$

the Weber-Hermite functions, i.e. the standard normalized eigenfunctions of the harmonic oscillator with frequency 1, then

$$\Psi_{\xi, m} = \sqrt[4]{B_c} \Omega_m\left(\sqrt{B_c} (y - \beta\xi)\right), \quad m \in \mathbb{Z}_+, \quad (8)$$

are the eigenfunctions of $\hat{H}_{\xi}$, and the corresponding eigenvalues are $B_c \left(\frac{1}{2} + m\right)$. Since the spectrum of $\hat{H}_{\xi}$ is independent of ξ it coincides with the spectrum of H as a set, and both are pure-point. On the other hand, since H is invariant in x , it has infinitely degenerate spectrum.

1.2 $\varepsilon_0 \neq 0 = \varepsilon_1$

Introducing the shifted variable $\tilde{y} = y - \beta\xi$, we can write the potential (5d) as

$$V_\xi(\tilde{y}) = \frac{B_c^2}{2} (\tilde{y} + \varepsilon_0 A^0(\tilde{y} + \beta\xi))^2. \quad (9)$$

Since A^0 is periodic and smooth (and therefore bounded), this is just a perturbation of the harmonic oscillator potential. $V_\xi(\tilde{y})$ will still tend to infinity as $|\tilde{y}|$ does, so that $\hat{H}_\xi$ has discrete spectrum. Some simple estimates using test functions show that for small ε_0 the eigenvalues will be within $C_m \varepsilon_0 \max |A^0|$ of the Landau levels, for some constants C_m (involving the maximum of $|y\Omega_m(y)|$) depending on m and B_c only; in particular, they are independent of ξ . A closer investigation shows that C_m is indeed bounded with respect to m .

Note that V_ξ is periodic in ξ with period $\frac{B_c}{2\pi} = \Phi$. Therefore, the spectrum of $H = \int_{\mathbb{R}}^{\oplus} \hat{H}_\xi d\xi$ consists of bands whose size is bounded by $2C_m \varepsilon_0 \max |A^0|$. These bands might degenerate into points.

1.3 $\varepsilon_0 \neq 0 \neq \varepsilon_1$

Now H is not independent of x any more. But at least it will be periodic in x with period 1, since A^1 is periodic. Using Fourier series on $L^2([0, 1]_x)$, we can decompose $L^2(\mathbb{R}^2)$ into a direct integral $\int_{[0, 1]}^{\oplus} L^2([0, 1]_x \times \mathbb{R}_y) d\xi$ such that

$$L^2(\mathbb{R}^2) \ni f \mapsto \hat{f}, \quad \hat{f}_\xi(x, y) = \sum_{l \in \mathbb{Z}} f(x + l, y) e^{-2\pi i \xi(x + l)}, \quad (10a)$$

$$f(x, y) = \int_{[0, 1]} \hat{f}_\xi(x, y) e^{2\pi i \xi x} d\xi; \quad (10b)$$

$$\begin{aligned} H = \int_{[0, 1]}^{\oplus} \hat{H}_\xi d\xi, \quad \hat{H}_\xi = \frac{1}{2} \left[\left(\frac{1}{i} \frac{\partial}{\partial x} + 2\pi\xi - B_c y - \varepsilon_0 A^0(y) \right)^2 + \right. \\ \left. + \left(\frac{1}{i} \frac{\partial}{\partial y} - \varepsilon_1 A^1(x, y) \right)^2 \right] \end{aligned} \quad (10c)$$

acting on functions periodic in x . Note that we keep denoting the fibre operator $\hat{H}_\xi$ for the new direct integral, in order to avoid an inflation of notation.

If we now choose the basis $(e^{2\pi i n x})_{n \in \mathbb{Z}}$ in $L^2([0, 1]_x)$, defining an isomorphism with $\ell^2(\mathbb{Z})$, and combine this with the isomorphism defining the direct integral above, we arrive at the direct integral $L^2(\mathbb{R}^2) = \int_{[0,1]}^{\oplus} \ell^2(\mathbb{Z}) \otimes L^2(\mathbb{R}) d\xi$ with

$$L^2(\mathbb{R}^2) \ni f \mapsto \hat{f}, \quad \hat{f}_{\xi,n}(y) = \int_{[0,1]} \sum_{l \in \mathbb{Z}} f(x+l, y) e^{-2\pi i [\xi l + (\xi+n)x]} dx, \quad (11a)$$

$$f(x, y) = \int_{[0,1]} \sum_{n \in \mathbb{Z}} \hat{f}_{\xi,n}(y) e^{2\pi i (\xi+n)x} d\xi; \quad (11b)$$

$$H = \int_{[0,1]}^{\oplus} \hat{H}_{\xi} d\xi, \quad \hat{H}_{\xi} = \frac{1}{2} \left[(2\pi(\nu + \xi) - B_c y - \varepsilon_0 A^0(y))^2 + \left(\frac{1}{i} \frac{d}{dy} - \varepsilon_1 \widehat{A^1}(y) \star \right)^2 \right] \quad (11c)$$

Here, ν is the operator of multiplication on $\ell^2(\mathbb{Z})$, i.e. $(\nu g)(n) = ng(n)$, and $\widehat{A^1}(y) \star$ is convolution with the Fourier series of $A^1(x, y)$ in x :

$$(\widehat{A^1}(y) \star g)(n) = \sum_{l \in \mathbb{Z}} g(l) \widehat{A^1}_{n-l}(y), \quad (12a)$$

$$\widehat{A^1}_n(y) = \int_{[0,1]} A^1(x, y) e^{-2\pi i n x} dx \quad (12b)$$

Note that, of course, our basis functions belong to the domain of the operator.

As a final step, we choose a special basis in $L^2(\mathbb{R})$, namely the eigenfunctions $(\Psi_{\xi,m})_{m \in \mathbb{Z}_+}$ described in subsection 1.1, equation (8). Thus we arrive at

Lemma 1. *There is a decomposition of $L^2(\mathbb{R}^2)$ into $\int_{[0,1]}^{\oplus} \ell^2(\mathbb{Z} \times \mathbb{Z}_+) d\xi$ with*

$$L^2(\mathbb{R}^2) \ni f \mapsto \hat{f}, \quad (13a)$$

$$\hat{f}_{\xi,n,m} = \int_{[0,1] \times \mathbb{R}} \sum_{l \in \mathbb{Z}} f(x+l, y) e^{-2\pi i [\xi l + (\xi+n)x]} \Psi_{\xi+n,m}(y) dx dy, \quad (13b)$$

$$f(x, y) = \int_{[0,1]} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}_+} \hat{f}_{\xi,n,m} e^{2\pi i (\xi+n)x} \Psi_{\xi+n,m}(y) d\xi \quad (13c)$$

such that H decomposes as

$$H = \int_{[0,1]}^{\oplus} \hat{H}_{\xi} d\xi, \quad (14a)$$

$$\left(\hat{H}_{\xi} \hat{f}_{\xi} \right)_{m,n} = B_c \left(\frac{1}{2} + m \right) \hat{f}_{\xi,m,n} \quad (14b)$$

$$+ \frac{1}{2} \varepsilon_0^2 \sum_{l \in \mathbb{Z}_+} \widetilde{A}_{l,m}^0(\beta(\xi + n)) \hat{f}_{\xi,l,n} \quad (14c)$$

$$- \varepsilon_0 \sum_{l \in \mathbb{Z}_+} A_{l,m}^0(\beta(\xi + n)) \hat{f}_{\xi,l,n} \quad (14d)$$

$$+ \frac{1}{2} \varepsilon_1^2 \sum_{l \in \mathbb{Z}_+} \sum_{k \in \mathbb{Z}} \widetilde{A}_{l,m}^{1(n-k)}(\beta(\xi + n)) \hat{f}_{\xi,l,k} \quad (14e)$$

$$- \varepsilon_1 \sum_{l \in \mathbb{Z}_+} \sum_{k \in \mathbb{Z}} \widehat{A}_{l,m}^{1(n-k)}(\beta(\xi + n)) \hat{f}_{\xi,l,k} \quad (14f)$$

where

$$\widetilde{A}_{l,m}^0(p) = \sqrt{B_c} \int_{\mathbb{R}} (A^0)^2(y+p) \Omega_l \left(\sqrt{B_c} y \right) \Omega_m \left(\sqrt{B_c} y \right) dy, \quad (15a)$$

$$A_{l,m}^0(p) = B_c^{\frac{3}{2}} \int_{\mathbb{R}} A^0(y+p) y \Omega_l \left(\sqrt{B_c} y \right) \Omega_m \left(\sqrt{B_c} y \right) dy, \quad (15b)$$

$$\widetilde{A}_{l,m}^{(k)}(p) = \sqrt{B_c} \int_{\mathbb{R}} \widehat{(A^1)^2}_k(y+p) \Omega_l \left(\sqrt{B_c} (y + \beta k) \right) \Omega_m \left(\sqrt{B_c} y \right) dy, \quad (15c)$$

$$\begin{aligned} \widehat{A}_{l,m}^{1(k)}(p) &= \frac{1}{l} B_c \int_{\mathbb{R}} \widehat{A^1}_k(y+p) \left[\Omega'_l \left(\sqrt{B_c} (y + \beta k) \right) \Omega_m \left(\sqrt{B_c} y \right) \right. \\ &\quad \left. - \Omega_l \left(\sqrt{B_c} (y + \beta k) \right) \Omega'_m \left(\sqrt{B_c} y \right) \right] dy. \end{aligned} \quad (15d)$$

As above, $\widehat{A^1}_k$ denotes the k -th Fourier coefficient of A^1 with respect to x , and analogously for $(A^1)^2$.

Proof. We have

$$\left(\hat{H}_{\xi} \hat{f}_{\xi} \right)_{m,n} = \sum_{l \in \mathbb{Z}_+} \sum_{k \in \mathbb{Z}} \hat{H}_{\xi,m,n;l,k} \hat{f}_{\xi,l,k}$$

so that we just have to compute the matrix elements in the given basis, for all the terms in (11c).

(14b) These are just the Landau eigenvalues in the case $\varepsilon_0 = 0 = \varepsilon_1$.

(14c) & (15a) For the terms with coefficient ε_0^2 we have to compute the matrix element of the square term $(A^0)^2$ which is

$$\sqrt{B_c} \int_{\mathbb{R}} (A^0)^2(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega_l \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) dy$$

(14d) & (15b) The terms with coefficient ε_0 in (11c) give the matrix element of the mixed term $(B_c y - 2\pi(\nu + \xi))A^0$, the calculation is the same as above.

(14e) & (15c) The coefficient ε_1^2 singles out the square term $\widehat{A^1} \star \widehat{A^1} \star = \widehat{(A^1)^2} \star$, i.e. convolution with the Fourier series of $(A^1)^2$. Its matrix element is

$$\sqrt{B_c} \int_{\mathbb{R}} \widehat{(A^1)^2}_{n-k}(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) dy$$

so that shifting y as above gives the desired result.

(14f) & (15d) The term with coefficient ε_1 is $\imath \left(\frac{d}{dy} \circ \widehat{A^1} \star + \widehat{A^1} \star \frac{d}{dy} \right)$. The first part can be written $\frac{d}{dy} \circ \widehat{A^1} \star = \widehat{A^1}' \star + \widehat{A^1} \star \frac{d}{dy}$ which has matrix element

$$\begin{aligned} & \sqrt{B_c} \int_{\mathbb{R}} \left[\widehat{A^1}'_{n-k}(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) \right. \\ & \quad \left. + \sqrt{B_c} \widehat{A^1}_{n-k}(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega'_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) \right] dy. \end{aligned}$$

On the other hand, we can use partial integration for the matrix element of the second part, which is $\widehat{A^1} \star \frac{d}{dy}$:

$$\begin{aligned} & B_c \int_{\mathbb{R}} \widehat{A^1}_{n-k}(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega'_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) dy \\ & = \sqrt{B_c} \int_{\mathbb{R}} \left[-\widehat{A^1}'_{n-k}(y) \Omega_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) \right. \\ & \quad \left. - \sqrt{B_c} \widehat{A^1}_{n-k}(y) \Omega'_m \left(\sqrt{B_c} (y - \beta(\xi + n)) \right) \Omega_l \left(\sqrt{B_c} (y - \beta(\xi + k)) \right) \right] dy. \end{aligned}$$

These two parts add up to the desired result.

□

Remark 1. Note that the functions $\widetilde{A}_{l,m}^0(p)$, $A_{l,m}^0(p)$, $\widetilde{A}_{l,m}^{(k)}(p)$, $\widehat{A}_{l,m}^{(k)}(p)$ have period 1 in p . Also, due to the decay of the Weber-Hermite functions $\Omega_m(y)$ in y , the functions $\widetilde{A}_{l,m}^{(k)}(p)$, $\widehat{A}_{l,m}^{(k)}(p)$ exhibit exponential decay in k .

Remark 2. If A^1 depends on x only then $\widetilde{A}_{l,m}^{(k)}(p) = \delta_{k,0} \delta_{l,m} \overline{(A^1)^2}$ with the average $\overline{(A^1)^2}$ of $(A^1)^2$ with respect to x . If we assume that $\overline{A^1} = 0$ (we can always add a constant to A^1 to achieve this, without changing the magnetic field) then $\widehat{A}_{l,m}^{(k)}(p) = 0$.

Remark 3. In the simplest non-trivial case $A^0(y) = \cos(2\pi y)$, using parity of the Weber-Hermite functions we get

$$A_{l,m}^0(p) = a_{l,m} \cos(2\pi p) + b_{l,m} \sin(2\pi p),$$

where $a_{l,m} = 0$ if $m + l$ is even, and $b_{l,m} = 0$ if $m + l$ is odd. Similarly,

$$\widetilde{A}_{l,m}^0(p) = \frac{1}{2} \delta_{l,m} + c_{l,m} \cos(4\pi p) + d_{l,m} \sin(4\pi p),$$

where $c_{l,m} = 0$ if $m + l$ is odd, and $d_{l,m} = 0$ if $m + l$ is even.

Remark 4. Creation and annihilation operators on harmonic oscillator functions yield the relations

$$\Omega'_m(y) = \sqrt{\frac{m}{2}} \Omega_{m-1} - \sqrt{\frac{m+1}{2}} \Omega_{m+1}, \quad (16a)$$

$$y \Omega_m(y) = \sqrt{\frac{m}{2}} \Omega_{m-1} + \sqrt{\frac{m+1}{2}} \Omega_{m+1}. \quad (16b)$$

Using these we can express $A_{l,m}^0$ and $\widehat{A}_{l,m}^{(k)}$ solely in terms of A^0 , A^1 and Weber-Hermite functions, without referring to their derivatives or multiplication by y .

2 Main results

Let $P_{\xi,m}^0$ denote the projection on the eigenspace of the m -th Landau level. Lemma 1 tells us that the action of $\hat{H}_\xi$ in this eigenspace, i.e. the part of $\hat{H}_\xi$

which is ‘diagonal in m ’, amounts to:

$$\begin{aligned} & \left(P_{\xi,m}^0 \hat{H}_\xi P_{\xi,m}^0 \right) \hat{f}_{\xi,m,n} \\ &= \left(B_c \left(\frac{1}{2} + m \right) + \frac{1}{2} \varepsilon_0^2 \widetilde{A}_{m,m}^0 (\beta(\xi + n)) - \varepsilon_0 A_{m,m}^0 (\beta(\xi + n)) \right) \hat{f}_{\xi,m,n} \quad (17) \\ &+ \sum_{k \in \mathbb{Z}} \left[\frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,m}^{1(n-k)} (\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,m}^{1(n-k)} (\beta(\xi + n)) \right] \hat{f}_{\xi,m,k} \end{aligned}$$

For fixed ξ and m this is a one-dimensional difference operator with quasiperiodic coefficients and exponentially decaying off-diagonal (i.e. $k \neq 0$) terms. If we choose A^1 to be independent of y as in Remark 2 there are no off-diagonal terms at all. If we furthermore choose $A^0(y) = \cos(2\pi y)$ as in Remark 3 then (17) will be similar to the Almost Matthieu operator, with a slightly more complicated potential. Indeed, if we look at terms of order up to ε_0 only it will be exactly the Almost Matthieu operator. In the case of constant magnetic field this observation goes back to Hofstadter (1976).

If ε_0 and ε_1 are small enough then H will have invariant subspaces E_m which are close to the eigenspaces of the Landau levels. We will construct a unitary transformation to achieve the following:

Theorem 1. *For every small enough ε_0 and ε_1 there is an $M(\varepsilon_0, \varepsilon_1)$ such that $M(\varepsilon_0, \varepsilon_1) \rightarrow \infty$ as both $\varepsilon_0, \varepsilon_1 \rightarrow \infty$, and such that for $m \leq M(\varepsilon_0, \varepsilon_1)$ the invariant subspace E_m of H and the restriction of H to E_m have a direct integral decomposition*

$$E_m = \int_{[0,1]}^{\oplus} \ell^2(\mathbb{Z}) d\xi, \quad (18a)$$

$$H_m := H|_{E_m} = \int_{[0,1]}^{\oplus} H_{m,\xi} d\xi. \quad (18b)$$

Furthermore, $H_{m,\xi}$ acts on $g \in \ell^2(\mathbb{Z})$ as a one-dimensional difference operator with exponentially decaying coefficients close to those of (17):

$$(H_{m,\xi} g)(n) = d_m(\beta(\xi + n)) g(n) + \sum_{k \in \mathbb{Z}} a_m(n - k, \beta(\xi + n)) g(k), \quad (19a)$$

$$\|d_m(\cdot) - [B_c(\frac{1}{2} + m) - \varepsilon_0 A_{m,m}^0(\cdot)]\|_{C^2(S^1)} < C_0 \varepsilon_0^2, \quad (19b)$$

$$\sum_{k \in \mathbb{Z} \setminus \{0\}} \|a_m(k, \cdot) e^{\delta|k|}\|_{C^2(S^1)} < C_1 \varepsilon_0 \quad (19c)$$

for some $\delta > 0$

We denote by $\|\cdot\|_{C^2(S^1)}$ the sum of the supremum norms of the derivatives of order 0, 1, 2.

As in (Dinaburg et al., 1997) we can now make use of the results of Dinaburg (1997) on ergodic families of operators. In order to apply these results we need the following assumptions:

diophantine There are $C > 0$, $\kappa > 0$ such that $|\{\beta n\}| > C/|n|^\kappa$ for all $n \in \mathbb{Z} \setminus 0$. Here, $\beta = \frac{2\pi}{B_c} = \frac{1}{\Phi}$ as before, and $\{\cdot\}$ denotes the fractional part.

smoothness A^0 and $A^1(x, y)$ are smooth; furthermore, all derivatives $\frac{\partial^j A^1}{\partial y^j}(x, y)$ are analytic in $|\Im x| < \delta$ for some common $\delta > 0$.

Morse $A_{m,m}^0$ is a Morse function on S^1 with exactly two critical points.

Theorem 2. *Let β be diophantine and A^0, A^1 smooth as defined above, and $M > 0$ such that $A_{m,m}^0$ is a Morse function with exactly two critical points for all $m \leq M$. Then there are $\tilde{\varepsilon}_0, \tilde{\varepsilon}_1 > 0$ depending on M such that for all $\varepsilon_0 < \tilde{\varepsilon}_0, \varepsilon_1 < \tilde{\varepsilon}_1$ the following are true for all $m \leq M$:*

1. *There are 1-periodic measurable functions λ_m such that for every $n \in \mathbb{Z}$ and almost every $\xi \in [0, 1]$, $\lambda_m(\beta(\xi + n))$ is an eigenvalue of $H_{m,\xi}$ and therefore in the spectrum of H_m . Furthermore,*

$$\|\lambda_m(\cdot) - [B_c(\frac{1}{2} + m) - \varepsilon_0 A_{m,m}^0(\cdot)]\|_{L^\infty(S^1)} < \text{const } \varepsilon_0^2. \quad (20)$$

2. *There are 1-periodic measurable functions $f_{m,l,k}, l \in \mathbb{Z}_+, n \in \mathbb{Z}$, decaying exponentially in n such that for almost every $\xi \in [0, 1]$,*

$$\sum_{\substack{l \in \mathbb{Z}_+ \\ k \in \mathbb{Z}}} |f_{m,l,n}(\beta\xi)| (l^2 + 1) e^{2\delta|n|} < \infty, \quad (21)$$

and for every $k \in \mathbb{Z}$, the series

$$\Phi_{m,\xi,k}(x, y) = \sum_{\substack{l \in \mathbb{Z}_+ \\ n \in \mathbb{Z}}} f_{m,l,n-k}(\beta(\xi + k)) e^{2\pi i(\xi+n)x} \Psi_{\xi+n,l}(y) \quad (22)$$

and all its derivatives converge uniformly in x, y . $\Phi_{m,\xi,k}$ is an eigenfunction of $H_{m,\xi}$ and therefore a generalized eigenfunction of H , with

eigenvalue $\lambda_m(\beta(\xi + k))$. Moreover, for every $N > 0$ and for $\varepsilon_0, \varepsilon_1$ small enough (depending on N),

$$|\Phi_{m,\xi,k}(x, y)| \leq \frac{\text{const}}{y^{2N} + 1} \quad (23)$$

with the constant depending on ξ, k, m, N .

3. H_m is uniformly ε_0 -close to band structure: H_m is unitarily equivalent to multiplication by the function $\lambda_m(\beta \cdot)$. The Lebesgue measure of the spectrum of H_m is $\varepsilon_0 |\text{ran } A_{m,m}^0| + O(\varepsilon_0^2)$.

Remark 5. One can reduce the smoothness requirements somewhat (to analyticity in x of a finite number of derivatives in y), thereby weakening the result on decay of the generalized eigenfunctions in y .

Remark 6. Of course one can also include an electric potential into the picture; Dinaburg et al. (1997) did so in the case of constant magnetic field, i.e. $A^0 \equiv A^1 \equiv 0$. The point in our work is that the magnetic field perturbation alone is strong enough to deform the Landau levels into a spectral set with positive measure.

3 Reduction

$\hat{H}_\xi$ is a double matrix operator on $\ell^2(\mathbb{Z}_+ \times \mathbb{Z})$ with indices $(m, n) \in \mathbb{Z}_+ \times \mathbb{Z}$. We decompose it as $\hat{H}_\xi = \mathcal{D}_1 + \mathcal{M}_1 + \mathcal{O}_1$, where $\mathcal{D}_1$ is diagonal in m and n , $\mathcal{O}_1$ is off-diagonal in m and contains only the first row and the first column, and $\mathcal{M}_1$ is the remainder. Note that both $\mathcal{O}_1$ and $\mathcal{M}_1$ are of order ε_0 (we always assume $\varepsilon_1 < \varepsilon_0$).

Our goal is to find a unitary transformation U which kills the terms in $\mathcal{O}$ (they represent the interaction between different Landau levels). U should leave the rest of the structure basically untouched. We will show how to accomplish this for the interaction between the 0-th and all other bands; extending this to M off-diagonal terms is a trivial generalisation.

The strategy is as follows: We define U as $U = \prod_{j \in \mathbb{N}} U_j$, where U_j eliminates off-diagonal terms up to (and including) order ε_0^j . Each transformation is of the form $U_j = e^{iW_j}$ for a Hermitian bounded W_j whose coefficients are

of order ε_0^j . We use the Baker-Hausdorff formula

$$e^{-\imath A} B e^{\imath A} = \sum_{r \in \mathbb{Z}_+} \frac{\imath^r}{r!} [B, A]_r, \quad (24a)$$

$$[B, A]_0 = B, \quad (24b)$$

$$[B, A]_{r+1} = [[B, A]_r, A] \quad (24c)$$

in order to find W_j . In fact, in the j -th step we will only have to consider the terms up to order ε_0^j in this formula, which are the terms $r = 0$ and $r = 1$. This gives us the equation

$$\mathcal{O}_1 + \imath [\mathcal{D}_1, W_1] = 0,$$

which reads as follows for the coefficients:

$$W_1(m, n; l, k) = 0 \quad \text{for } m > 0, \quad (25a)$$

$$\mathcal{O}_1(0, n; l, k) + \imath [\mathcal{D}_1(0, n; 0, n) - \mathcal{D}_1(l, k; l, k)] W_1(0, n; l, k) = 0 \quad (25b)$$

Because of

$$\begin{aligned} & |\mathcal{D}_1(0, n; 0, n) - \mathcal{D}_1(l, k; l, k)| > \\ & B_c l - \frac{1}{2} \varepsilon_0^2 \left(\left\| \widetilde{A}_{0,0}^0 \right\|_\infty + \left\| \widetilde{A}_{l,l}^0 \right\|_\infty \right) - \varepsilon_0 \left(\left\| A_{0,0}^0 \right\|_\infty + \left\| A_{l,l}^0 \right\|_\infty \right) \\ & - \frac{1}{2} \varepsilon_1^2 \left(\left\| \widetilde{A}_{0,0}^{(0)} \right\|_\infty + \left\| \widetilde{A}_{l,l}^{(0)} \right\|_\infty \right) - \varepsilon_1 \left(\left\| \widehat{A}_{0,0}^{(0)} \right\|_\infty + \left\| \widehat{A}_{l,l}^{(0)} \right\|_\infty \right) \\ & > \frac{1}{2} B_c l \end{aligned}$$

we can choose ε_0 small enough so that for all $l > 0$ there is no small denominator problem when solving equation (25b) for W_1 .

Defining $H_2 = U_1^* H U_1$ and repeating the above steps we arrive at the following:

Lemma 2. *We define W_j inductively by $[\mathcal{D}_j, W_j] = \imath \mathcal{O}_j$, and furthermore $U_j = e^{\imath W_j}$, $H_{j+1} = U_j^* H_j U_j$. Then we have:*

closeness to diagonal operator

$$\left\| \mathcal{D}_j(m, n; m, n) - \left[B_c \left(\frac{1}{2} + m \right) - \varepsilon_0 A_{m,m}^0 \right] \right\|_{C^2(S^1)} < \varepsilon_0 \delta_j \quad (26)$$

off-diagonal smallness

$$\sum_{l \in \mathbb{Z}_+} \|\mathcal{O}_j(0, n; l, n)\|_{C^2(S^1)} (l^s + 1) < \gamma_j \quad (27a)$$

$$\sum_{k \in \mathbb{Z} \setminus \{n\}} \sum_{l \in \mathbb{Z}_+} \|\mathcal{O}_j(0, n; l, k)\|_{C^2(S^1)} e^{2\delta|n-k|} (l^s + 1) < \varepsilon_1 \gamma_j \quad (27b)$$

smallness of mixed terms

$$\sum_{l \in \mathbb{Z}_+} \|\mathcal{M}_j(m, n; l, n)\|_{C^2(S^1)} (l^s + 1) < (m^{s+1} + 1) \delta_j \quad (28a)$$

$$\sum_{k \in \mathbb{Z} \setminus \{n\}} \sum_{l \in \mathbb{Z}_+} \|\mathcal{M}_j(m, n; l, k)\|_{C^2(S^1)} e^{2\delta|n-k|} (l^s + 1) < (m^{s+1} + 1) \varepsilon_1 \delta_j \quad (28b)$$

Furthermore, we have the relations

$$\gamma_{j+1} = \text{const } \gamma_j \delta_j, \quad (29a)$$

$$\delta_{j+1} = \delta_j (1 + \text{const } \gamma_j), \quad (29b)$$

$$W_j(0, n; l, k) \sim \frac{\mathcal{O}_j(0, n; l, k)}{l+1} \text{ as } l \rightarrow \infty, \quad (29c)$$

where the constants depend on s .

Later we will see that

$$\begin{aligned} \delta_j &< \text{const}_s \varepsilon_0 \\ \gamma_j &< (\text{const}_s \varepsilon_0)^j \end{aligned}$$

4 Estimates

In order to prove Lemma 2 we only need to check the induction hypothesis. For $j = 1$ we have the following non-vanishing matrix elements:

$$\begin{aligned} \mathcal{D}_1(m, n; m, n) &= B_c \left(\frac{1}{2} + m \right) + \frac{1}{2} \varepsilon_0^2 \widetilde{A}_{m,m}^0(\beta(\xi + n)) - \varepsilon_0 A_{m,m}^0(\beta(\xi + n)) \\ &\quad + \frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,m}^{(0)}(\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,m}^{(0)}(\beta(\xi + n)); \end{aligned} \quad (30a)$$

$$\mathcal{M}_1(m, n; l, k) \stackrel{n \neq k}{=} \frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,l}^{(n-k)}(\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,l}^{(n-k)}(\beta(\xi + n)), \quad (30b)$$

$$\begin{aligned} \mathcal{M}_1(m, n; l, n) &= \frac{1}{2} \varepsilon_0^2 \widetilde{A}_{m,l}^0(\beta(\xi + n)) - \varepsilon_0 A_{m,l}^0(\beta(\xi + n)) \\ &\quad + \frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,l}^{(0)}(\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,l}^{(0)}(\beta(\xi + n)); \end{aligned} \quad (30c)$$

$$\begin{aligned} \mathcal{O}_1(m, n; 0, n) &= \frac{1}{2} \varepsilon_0^2 \widetilde{A}_{m,0}^0(\beta(\xi + n)) - \varepsilon_0 A_{m,0}^0(\beta(\xi + n)) \\ &\quad + \frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,0}^{(0)}(\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,0}^{(0)}(\beta(\xi + n)), \end{aligned} \quad (30d)$$

$$\mathcal{O}_1(m, n; 0, k) \stackrel{n \neq k}{=} \frac{1}{2} \varepsilon_1^2 \widetilde{A}_{m,0}^{(n-k)}(\beta(\xi + n)) - \varepsilon_1 \widehat{A}_{m,0}^{(n-k)}(\beta(\xi + n)) \quad (30e)$$

Note that we may assume $\widehat{A}_{m,l}^{(0)} = 0$; we can always achieve this by changing A^1 by a function of y only, this does not change B . $\widetilde{A}_{m,l}^{(0)} = 0$ cannot be made disappear in the same way because it involves the average of $(A^1)^2$ instead of A^1 .

If we rewrite the induction hypotheses (26)-(29c) in terms of the coefficients of the vector potential we get:

$$\begin{aligned} \sum_{l \in \mathbb{Z}_+} \left\| \frac{1}{2} \varepsilon_0 \widetilde{A}_{m,l}^0(\beta(\xi + n)) - A_{m,l}^0(\beta(\xi + n)) \right. \\ \left. + \frac{1}{2} \varepsilon_1 \widetilde{A}_{m,l}^{(0)}(\beta(\xi + n)) - \widehat{A}_{m,l}^{(0)}(\beta(\xi + n)) \right\| < \text{const} (m^{s+1} + 1) \end{aligned} \quad (31a)$$

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \sum_{l \in \mathbb{Z}_+} \left\| \frac{1}{2} \varepsilon_1 \widetilde{A}_{m,l}^{(n-k)}(\beta(\xi + n)) - \widehat{A}_{m,l}^{(n-k)}(\beta(\xi + n)) \right\| e^{2\delta|n-k|} \\ < \text{const} (m^{s+1} + 1) \end{aligned} \quad (31b)$$

Since the vector potentials A^0, A^1 are periodic in y , equations (31a) and (31b) can be derived from the smoothness (analyticity) assumption and estimates

for

$$\int_{\mathbb{R}} e^{i\eta y} \Omega_m(y) \Omega_l(y) dy$$

in η . For this we can use the estimates from the case of constant magnetic field (Dinaburg et al., 1997, lemmata 6-8). Because of our Remark 4 there are no new estimates to prove.

5 Finishing the Proof

The columns of the transformation U define new basis vectors $b_{\xi,n,m}, n \in \mathbb{Z}, m \in \mathbb{Z}_+$. From the induction hypothesis (27b) we get

$$|b_{\xi,n,m}|(m^{N+1} + 1)e^{2\delta|n-j|} < \text{const}. \quad (32)$$

This implies that for each m , we can apply the main results of Dinaburg (1997) to the operator H_m , which is an ergodic family (in ξ) of difference operators with exponentially decaying almost periodic coefficients. As a consequence we get part 1 and the eigenfunction expansion in part 2, equations (21) and (22), of Theorem 2.

Uniform convergence of (22) and its derivatives follows from (32) and the relation (16a) for the derivatives. As in (Dinaburg et al., 1997), (32) also gives the estimate on the rate of decay in y , (23).

Finally, from part 1 we know that, for almost every ξ , the spectrum of $H_{m,\xi}$ is given by $(\lambda_m(\beta(\xi + n)))_{n \in \mathbb{Z}}$. If $E_{\xi,m,n}$ are the corresponding eigenspaces, then $E_{m,n} := \int^{\oplus} E_{\xi,m,n} d\xi$ are invariant subspaces such that the restriction of H to $E_{m,n}$ is unitarily equivalent to multiplication with the function $\xi \mapsto \lambda_m(\beta(\xi + n))$. On the other hand, $H_m = \bigoplus_{n \in \mathbb{Z}} E_{m,n}$. Therefore, since $H_m = \int^{\oplus} H_{m,\xi} d\xi$, the operator H_m is unitarily equivalent to multiplication with the function λ_m .

This together with the quantitative estimate (20) finally gives the assertion in part 3 about the measure of the spectrum.

References

- BIRMAN, M. S. & SUSLINA, T. A. (1998). Two-dimensional periodic magnetic Hamiltonian is absolutely continuous. *St. Petersburg Math. J.* **9**, no. 1, 21–32. English Translation of Algebra Anal.

- DINABURG, E. I. (1997). Some problems in the spectral theory of discrete operators with quasiperiodic coefficients. *Russian Math. Surveys* **52**, no. 3, 451–499. English Translation of Uspekhi Mat. Nauk
- DINABURG, E. I., SINAI, Y. G. & SOSHIKOV, A. B. (1997). Splitting of the low Landau levels into a set of positive measure under small periodic perturbation. *Comm. Math. Phys.* **189**, no. 2, 559–575
- GRUBER, M. J. (2001). Non-commutative Bloch theory. *J. Math. Phys.* **42**, no. 6, 2438–2465. math-ph/0006021
- HELFFER, B. & SJÖSTRAND, J. (1989). Semiclassical analysis for Harper’s equation III. Cantor structure of the spectrum. *Mém. Soc. Math. France (N.S.)* **39**, 1–124
- HOFSTADTER, D. R. (1976). Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields. *Phys. Rev. B (3)* **14**, 2239
- SOBOLEV, A. V. (1999). Absolute continuity of the periodic magnetic Schrödinger operator. *Invent. Math.* **137**, 85–112